

Application No. : 09/640,041  
Docket No. : PP-01615.002/200130.503

VERSION WITH MARKINGS TO SHOW CHANGES MADE

In the Specification:

Paragraph beginning at page 12, line 11, has been amended as follows:

Fusion proteins comprising EGFH2 or a biologically active or antigenic fragment thereof can be produced using methods known in the art. Such fusion proteins can be used therapeutically or can be produced in order to simplify the isolation and purification procedures. Histidine residues can be incorporated to allow immobilized metal affinity chromatography purification. Residues EQKLISEEDL (SEQ ID NO: 5) contain the antigenic determinant recognized by the myc monoclonal antibody and can be incorporated to allow myc monoclonal antibody-based affinity purification. A thrombin cleavage site can be incorporated to allow cleavage of the molecule at a chosen site; a preferred thrombin cleavage site is residues LVPRG (SEQ ID NO: 6). Purification of the molecule can be facilitated by incorporating a sequence, such as residues SAWRHPQFGG (SEQ ID NO: 7), which binds to paramagnetic streptavidin beads. Such embodiments are described in WO 97/25345, which is incorporated by reference.

JEP:mls  
701 Fifth Avenue, Suite 6300  
Seattle, Washington 98104-7092  
Phone: (206) 622-4900  
Fax: (206) 682-6031